

STUDY CONCERNING THE HONEY QUALITIES IN TRANSYLVANIA REGION

Maria Popa¹
Mihaela Vica²
Roxana Axinte³
Mirel Glevitzky⁴
Simona Varvara⁵

ABSTRACT: The sources of micro-organisms (yeasts and fungi) found in honey are nectar and pollen, honey processing areas, equipments that have not been properly cleaned or wrappings.

*There are few types of yeast in honey and the most common are *Saccharomyces melis*, which grows in media with water content over 20-25% and *Saccharomyces rosei*, which can ferment in media with 60% carbohydrates. Yeasts can produce microbiological faults in honey with more than 10² cells /g honey, stored at temperatures over 15 °C (Şindilar, E., 2000).*

Fungi can come from dust contamination, from the water with which installations or containers are washed and to a smaller degree, they can come from the honeybees. If they are found in honey in a vegetative state, they can metabolise carbohydrates, amino-acids and even pollen, causing various organoleptic changes (taste and smell of mildew).

The present paper is a comparative microbiological and physical-chemical analysis of various types of honey (polyfloral, tilia, acacia, sunflower, and honeydew) collected from beekeepers. The results have enabled us to make correlations between moisture, acidity, pH and the microbiological characteristics of the tested honey samples and processors.

Key words: quality, product quality, honey quality

JEL codes: M31

Introduction: Aims And Background

The system of ensuring food products in sufficient amounts is on a continuous descending line. At the present time, the integrity, quality, sanitation and nutritional value of food products are paid increased attention.[1]. The concept of “food safety based on the general hygiene principles of food products and on the HACCP method adopted by *Codex Alimentarius* comes to reduce or control biological, physical and/or chemical contaminations [2]. Maintaining the contamination level within accessible, minimum limits leads to a linear and constant process when it comes to ensuring quality and food safety. Thus, the harvesting, collecting, manufacturing and storing of honey are a field in a continuous evolution and adjustment to the European legislation, industry needs and the harsh market competition.

Honey represents a semi liquid, yellow, sweet and flavoured foodstuff, with a great biological and caloric value (it contains sugar, vitamins and enzymes), collected and produced by the bees from nectar, manna or sweet juice that can be found in different parts of plants and trees. Being a “noble” product of the bee, honey has a wide range of action depending on its kind [3].

¹“1 Decembrie 1918” University of Alba Iulia, Dept. of Topography, e-mail: mariapopa2010@yahoo.com

²“1 Decembrie 1918” University of Alba Iulia, Dept. of Topography

³ “1 Decembrie 1918” University of Alba Iulia, Dept. of Topography

⁴ Veterinary Health County Department Alba

⁵ “1 Decembrie 1918” University of Alba Iulia, Dept. of Topography

The physical-chemical and hygienic qualities of honey constitute indicators that offer information regarding the energetic and nutritional quality, as well as the possibility of falsifying honey. The authentication of honey is assessed through its physical-chemical parameters falling within the limits imposed by the present legislation. Falsifying agents or inappropriate thermal treatments are identified through the HMF (HydroxyMethylFurfuraldehyde) content.

The microorganisms in honey come from nectar and pollen, from the processing area, from the insufficiently washed machines or containers. The more frequently encountered sporulated microorganisms belong to the *Bacillus* type. The non-sporulated bacteria (*Micrococcus*, *Pseudomonas*, *Flavobacterium*) are less numerous, coming from the floral organs or the digestive tract of the bees. The filamentous fungi, being more spread in nature and having thermal resistant spores, with a great capacity of surviving, can be introduced in honey even by man, through dust, through the water installations or containers or even by the bees through pollen [4]. The microbes found in honey are not dangerous for the consumers' health. Even if *Aspergillus flavus* is found, there are no favourable conditions for aflatoxin.

Yeasts are present in a small number in normal honey. They are taken by the bees into the hive, but in this case they do not decrease the quality of honey. They can be found in larger amounts due to extraction and manipulation areas, installations, etc. and are especially dangerous through their number and acid. Due to their chemical composition, honey favours its fermentation, frosting and crystallisation [5].

The presence of microorganisms in honey can sometimes influence the stability of the product and its hygienic quality. Normal honey must lack pathogenic microorganisms or microorganisms that produce enteric illnesses.

Honey was often incriminated as a source of *Clostridium botulinum* spores responsible for causing infant botulism [6]. Huttanem and col. (1981), in 80 collected samples from the hives spores of *Clostridium* were not found. In another study, Kauttera and col (1982), out of 100 honey samples they found only two contaminated with *Clostridium botulinum* spores. In a study on honey that took place in Portugal a low percentage of contamination with *B.cereus* and fungi was found: yeast, *Mucor* spp., *Penicillium* spp and a few species of the *Aspergillus* type, particularly *Asp. flavus*, *Asp. candidus*, *Asp. fumigatus* and *Asp. niger*. In another study done on 80 honey samples, there were no *Clostridium perfringens* spores detected in any sample, *B. cereus* was identified in 11 samples (13.7%). Yeast and mould were detected in 71 samples (88.8%). There were three types of yeast that were identified (*Aspergillus*, *Penicillium* and *Mucor*) and two types of yeast (*Saccharomices* and *Candida*). The predominant species of *Aspergillus* were *A. flavus*, *A. niger*, *A. fumigatus* and *A. candidus*. No sample presented aflatoxin contamination [7]

Microbiological contamination during or after processing honey was demonstrated by the absence of the microorganisms in the samples collected from primary sources and by the presence of a certain type of bacterium (*Bacillus* spp) and eight types of fungi (more frequently *Candida*, *Aspergillus*, *Geotrichum* and *Rhizopus*) in the collected samples on local markets. This fact indicates the contamination from secondary sources during manipulations and previous processes. The contamination with fungi and bacteria indicate inadequate hygiene conditions during collecting, manipulating, processing and storing [8].

Experimental

Twenty bulk liquid honey samples of known origin, aseptically collected from beekeepers located in different areas of Transylvania (Romania) during 2008 and presented in table 1, were used for analysis. Each honey sample was purchased in duplicate in sterilised sealed jars of 200 g.

The main **physical and chemical indicators** (hydroxymethylfurfuraldehyde (HMF), humidity, acidity and pH) that reflect the honey quality were determined according to the methods proposed in the Harmonized Methods of International Honey Commission [9].

The HMF content was determined according to the White method using a UV–VIS spectrophotometer (model T80 PG Instruments, UK).

The absorbance of the honey samples was determined against the reference solution at 284 nm and 336 nm. The HMF content expressed in mg/kg honey was calculated according to the following equation:

$$\text{HMF} = (A_{284} - A_{336}) * 149.7 * 5 * D/W \text{ [mg/kg]} \quad (1)$$

where:

A_{284} = absorbance at 284 nm

A_{336} = absorbance at 336 nm

$$149.7 = \frac{126 \times 1000 \times 1000}{16830 \times 10 \times 5} = \text{Constant} \quad (2)$$

D = dilution factor (in case when dilution is necessary)

W = weight of honey sample (g)

Moisture content was determined measuring the refractive indices at 20⁰C by an ABBE refractometer. The corresponding moisture content was calculated from the refractive index of the honey by reference to a standard table.

The acidity of honey is the content of all free acids, expressed in milliequivalents/kg honey.

The honey sample (10 g) was dissolved in 75 ml carbon dioxide-free water and the pH value was measured using a pH-meter (Inolab level 2, WTW). The same solution was titrated with 0.1 M NaOH solution to pH = 8.30, using an automatic titrator (Titroline Alpha Plus, Schott Instruments).

Free acidity, express as milliequivalents or millimoles acid/ kg honey = ml of 0.1 M NaOH x 10.

From a **microbiological** point of view, the contamination of the samples was done by determining the total number of aerobic mesophilic bacteria (NTG) and determining the yeasts and moulds. The used diluting liquid was peptonate physiological serum: 10g of sample were homogenised with 90 ml of SFP, obtaining the diluted solution 10⁻¹.

Total number of germs

Petrifilm Aerobic Count Plate is used –3M Microbiology producer USA.

Petrifilm is a reactive film covered with a dehydrated culture medium which contains standard nutrients, a jelly making agent which is soluble in cold water and a tetrazolium indicator which facilitates the enumeration of colonies.

For each sample, two Petrifilms are used, placing them on a flat surface. The upper part of the film is lifted and, with a sterile dropper we put 1 ml at a time from the diluted solution 10⁻¹ on each of the two slates, the upper film is placed on the sample and is distributed with the help of the applying tool, and then some time is given to let it solidify for at least one minute. The Petrifilms are incubated at 30°C ± 1°C for 72h. All the red colonies are counted, regardless of the size or intensity.

Yeast and mould]

Take two sterile boxes of Petri. With a sterile dropper pour 1 ml of the 10⁻¹ diluted solution in every box. About 15 ml of yeast-glucose-cloramfenicol-agar extract is poured (Organics producer), previously melted and maintained at 45⁰±1⁰C in water, in every Petri box. Everything is carefully mixed, it is left to solidify, placing the Petri boxes on a horizontal, cold surface. A witness box is prepared, with 15 ml medium, in order to verify the sterility. The boxes are placed with the lid down, in the incubator, 25⁰±1⁰C. After three, four and five days of incubation, the colonies from each Petri box are counted. Evaluation: 100 CFU/g.

In order to identify the species and capture the images, a Hund Wetzlar H600LL microscope connected to a PC was used, using the Pinuacle TV Centre programme.

Results and discussion

Table 1 shows the physical-chemical indices: moisture content, pH, acidity and HMF of analyzed samples.

Table no. 1

The results of the physical-chemical analyses of the honey samples

Sample no.	Sample code	Moisture content (%)	pH	Acidity (meq /kg)	HMF (mg/kg)
1.	Polyfloral honey	15.6	3.72	41	12.2
2.	Polyfloral honey	16.8	3.99	23	52.3
3.	Linden honey	16.4	4.39	16	14.9
4.	Linden honey	17	4.86	11	1.0
5.	Acacia honey	18	3.75	12	4.3
6.	Acacia honey	19.4	2.89	14	8.4
7.	Forest honey	16.2	3.66	28.9	5.7
8.	Forest honey	15	3.98	43.8	21.7
9.	Sun flower honey	16.4	3.67	22.6	22.3
10.	Sun flower honey	19.8	3.59	20.9	23.8
11.	Polyfloral honey	16.4	3.6	37.3	31.8
12.	Polyfloral honey	15.4	4.01	20.6	7.4
13.	Polyfloral honey	15.6	3.9	55.6	4.5
14.	Polyfloral honey	16.4	3.78	28.5	2.3
15.	Polyfloral honey	16.6	4.04	30.4	8.5
16.	Polyfloral honey	14.6	3.8	24.4	30.0
17.	Linden honey	16.4	4.36	20.0	24.8
18.	Linden honey	16.4	4.01	23.4	14.2
19.	Acacia honey	16.4	3.9	14.3	64.3
20.	Acacia honey	16.8	3.9	14	10.7

The physical-chemical parameters values were within the reference ranges presented in UE or national reglementation. Exception is sample number 2 and 19 for HMF value and sample 13 for acidity level.

In order to establish the microbiological characteristics of the honey samples, the parameters from table 2 were analysed.

Table no. 2

The microbiological control of the honey samples

Number of the sample	NTG/g	DM/g
1	30	30 (<i>Penicillium</i> spp.)
2	95	15 (<i>Rhizopus</i> spp., <i>Aspergillus</i> spp., <i>Penicillium</i> spp.)
3	45	<10
4	15	15 (<i>Absidia</i> spp.)
5	20	<10
6	20	20 (<i>Penicillium</i> spp.)
7	40	40 (<i>Penicillium</i> spp., <i>Aspergillus</i> spp.)
8	45	20 (<i>Penicillium</i> spp., <i>Aspergillus</i> spp.)
9	35	<10
10	10	10 (<i>Fusarium</i> spp.)
11	30	<10
12	25	<10
13	<10	<10
14	20	10 <i>Aspergillus</i> spp
15	45	10 (<i>Penicillium</i> spp.)
16	10	<10
17	20	<10

18	25	<10
19	<10	<10
20	20	<10

The antimicrobial character of honey is confirmed by the results regarding the TNG (the total number of germs): in all the analysed samples it is under 100 CFU/g – value settled by the present legislation- which represents a reduced contamination with aerobic mesophilic germs. Out of the 20 analysed samples, the greatest microbiological importance was represented by sample no. 2 with 95 CFU/g.

Regarding the yeasts and moulds, we notice that there is no yeast in all the 10 analysed samples and the number of mould does not exceed 40/g in any sample. The most frequently encountered one is the *Penicillium* type (in five samples), then *Aspergillus* (in three samples). In each sample the fungi from the *Absidia* (*Mycocladius*), *Rhizopus* and *Fusarium* types were determined. We can notice that in the polyfloral honey sample from the *Penicillium*, *Aspergillus* and *Rhizopus* types can be found. Also, fungi of the *Penicillium* type were identified in the acacia honey sample.



Figure no. 1 - The *Penicillium* type
(Samples 1,2,6,7,8,15)




Figure no. 2 - The *Rhizopus* type
(Sample 2)




Figure no. 3 - The *Aspergillus* type
(Sample 2,7,8,14)




Figure no. 4 - The *Absidia* type
(Sample 4)




Figure no. 5 - The *Fusarium* type
(Sample 10)

We can notice the presence of TNG in the samples subjected to various handling and also a variety regarding the types of fungi encountered in the honey samples. This fact confirms a contamination from a microbiological point of view during its manipulation by the beekeepers and the primary honey treatments, which indicate unsatisfying hygiene conditions.

Figures 6 and 7 present the results of the correlated log of bacterial number and moisture separately for polyfloral honey samples from other samples and except the two samples (13 and 19), where the microbial growth is lower than 10.




Figure no. 6 - Correlation between total counts in polyfloral honey sample and their moisture content




Figure no. 7 - Correlation between total counts in linden, acacia, forest and sun flower honey sample and their moisture content

Slope of regression line log of bacterial number over moisture is $y = 0,6976x - 7,7593$, $r^2 = 0,6378$ in case of polyfloral honey samples and $y = -0,253x + 7,5075$, $r^2 = 0,5714$ for linden, acacia, forest and sun flower honey sample.

The dependences between TNG and the water content from the analysed samples were emphasized. The correlative analysis shows that there are appreciating correlations between the microbiota and the physical-chemical honey parameters.

Conclusions

The study allowed the qualitative analysis of the honey samples collected from beekeepers in Transylvania.

The experimental values of the physical-chemical and microbiological parameters of honey demonstrate the following:

- The presence of mould (especially of mould) in the *Penicillium*, *Aspergillus*, *Absidia*, *Rhizopus*, *Fusarium* types, but which cannot exceed the limit values. These facts, as well as the favourable conditions can lead to generating and developing micotoxins.
- Contamination from secondary sources during the manipulations due to the inadequate hygiene conditions during the selection, manipulation and storing.
- The importance of corresponding processing of honey: filtration, dehydration, liquefaction, pasteurization (70-78⁰C for 5-6 min.), cooling (sudden at 42⁰C) and wrapping in order to stop or destroy the present microorganisms.
- The physical-chemical parameters were within the limits imposed by the present legislation, except for 2.13 and 19 samples.
- Correlating the physical-chemical and microbiological results is necessary in order to check the quality and the sanitation of honey. This fact constitutes practical proof in ensuring food safety.

References:

1. Amon, S. S., Damus, K. Si Chin, 1981. J. Infant botulism: epydemiology and relation to sudden infant death syndrome. *Epidemiol. Rev.* 345-66.
2. Gillian, M, Moffett, J. O. si Kauffeld, N. M., 1983. Examination of floral nectar of citrus, cotton and Arizona desert plants for microbes. *Apidologie*, 299-302.
3. Herminia Martina Martins, M. Ligia Martins, Fernando M. A. Bernardo, 2003. Bacillaceae spores, fungi and aflatoxins determination in honey. *Revista Portuguesa de Ciencias Veterinarias*, 989 (546), 85-88.
4. Mărghitaş Liviu Alexandru, 2008. Bees and their products, Ceres Publishing, Bucharest, p. 280.
5. Savu, Constantin, 2008. Hygiene and product control. Semne Publishing, Bucureşti.
6. Stefan Bogdanov, 2002. Harmonised methods of the international honey commission. 1-62.
7. Tchoumboue Joseph, Awah-Ndukum Julius, Foneth Florence A, Dongock N Delphine, Pinta Jonnas and Mvondo Ze Antoine, 2007. Physico-chemical and microbiological characteristics of honey from the sudano-guinean zone of West Cameroon. *African Journal of Biotechnology* Vol 6(7), pp 908-913.
8. Tofan, Clemansa, 2004. Food Microbiology. AGIR Publishing, Bucharest, p. 285.
9. *** 2007. National food safety best practice guidelines. Uranus Publishing, Bucharest, p.14.
10. ***SR ISO 7954-2001 General directives for the count of yeasts and moulds. The technique of colnies counting at 25⁰C.